

**SECTION NINE:**

**REFERENCES CITED**

- Abernethy, B. and Rutherford, I.D., 1998. Where along a river's length will vegetation most effectively stabilize stream banks? *Geomorphology*, v. 23, p. 55-75.
- Anon., 1998. Draft strategic plan for restoration of endangered coho salmon south of San Francisco Bay: California Department of Fish and Game Open File report.
- Barbour, M.T., Gerritsen, J., Snyder, B.D., and Stribling, J.B., 1999. Revision to rapid bioassessment protocols for use in stream and rivers: periphyton, BMIs and fish. EPA 841-D-97-002. U.S. Environmental Protection Agency. Washington DC.
- Bash, J., Berman, C., and Bolton, S., 2001. Effects of turbidity and suspended solids on salmonids. Report by Center for Streamside Studies, University of Washington prepared for the Washington Department of Transportation. 74 pp.  
<http://depts.washington.edu/cssuw/Publications/Salmon%20and%20Turbidity.pdf> (June 2003).
- Bell, M.C., 1986. Fisheries handbook of engineering requirements and biological criteria. U.S. Army Corps of Engineers, Office of the Chief of Engineers, Fish Passage Development and Evaluation Program, Portland, Oregon.
- Benda, L.E., Bigelow, P., and Worsley, T.M., 2002. Recruitment of wood to streams in old-growth and second-growth redwood forests, northern California, USA. *Canadian Journal of Forestry Research*, v. 32, p. 1460-1477.
- Berg, L. and Northcote, T.G., 1985. Changes in territorial, gill-flaring and feeding behavior in juvenile coho salmon (*Oncorhynchus kisutch*) following short term pulses of suspended sediment. *Canadian Journal of Fisheries and Aquatic Sciences*, v. 42, p. 1410-1417.
- Bisson, P.A. and Bilby, R.E., 1982. Avoidance of suspended sediment by juvenile coho salmon. *North American Journal of Fisheries Management*, v. 4, p. 371-374.
- Bjornn, T.C., 1978. Survival, production, and yield of trout and chinook salmon in the Lemhi River, Idaho. Idaho Department of Fish and Game. Federal Aid to Fish Restoration Project F-49-R. 57 pp.
- Bjornn, T.C. and Reiser, D.W., 1991. Habitat requirements of salmonids in streams. In: Meehan, W.R. (ed). Influences of forest and rangeland management on salmonid fishes and their habitat. American Society Special Publication. p. 83-138.
- Brabb, E.E. and Pampeyan, E.H., 1972. Preliminary map of landslides in San Mateo County, California. U.S. Geological Survey Miscellaneous Field Studies Map MF-344, scale 1:62,500.
- Brabb, E.E., Pampeyan, E.H., and Bonilla, M.G., 1972. Landslide susceptibility in San Mateo County, California. U.S. Geological Survey Miscellaneous Field Studies Map MF-360, scale 1:62,500.
- Brabb, E.E. and Olson, J.A., 1986. Map showing faults and earthquake epicenters in San Mateo County, California. U.S. Geological Survey Miscellaneous Investigations Map I-1257-F, scale 1:62,500.
- Brabb, E.E., Graymer, R.W., and Jones, D.L., 2000. Geologic map and map database of the Palo Alto 30' x 60' quadrangle, California. U.S. Geological Survey Miscellaneous Field Studies Map MF-2332, scale 1:100,000.

- Bunte, K. and Abt, S.R., 2001. Sampling frame for improving pebble counts accuracy in coarse gravel-bed streams. *The Journal of the American Water Resources Association*, v. 37, p. 1001-1014.
- California Data Exchange Center, 2003. Data obtained from the California Data Exchange Center website. [www.cdec.water.ca.gov](http://www.cdec.water.ca.gov) (April 2003).
- California Department of Fish and Game, 2002. Culvert criteria for fish passage. <http://www.dfg.ca.gov/nafwb/pubs.html> (December 2003).
- CALWATER, 1997. Teale GIS Solutions Group. <http://www.ca.nrcs.usda.gov/features/calwater/>
- Cannon, S.H. and Ellen, S., 1985. Rainfall conditions for abundant debris avalanches, San Francisco Bay region, California. *California Geology*, v. 38, p. 267-272.
- Church, M.A., McLean, D.G., and Wolcott, J.F., 1987. Riverbed gravels; sampling and analysis. In: Thorne, C.R., Bathurst, J.C., and Hey, R.D., (eds). *Sediment transport in gravel-bed rivers*. John Wiley and Sons, Chichester, United Kingdom. 1012 pp.
- Clahan, K.B., Hall, N.T., and Wright, R.H., 1994. Late Holocene slip rate and slip events for the San Francisco Peninsula segment of the San Andreas fault. *American Geophysical Union Abstract*.
- Clark, J.C., 1970. Geologic map of the southwestern Santa Cruz Mountains between Ano Nuevo Point and Davenport, California. U.S. Geological Survey Open File Report, scale 1:24,000.
- Collins, L., 2001. Wildcat Creek watershed: A scientific study of physical processes and land use effects. Report prepared for Contra Costa Clean Water Program. San Francisco Estuary Institute, Oakland California. June 2001. 85pp.
- Collins, L. and Collins, J., 1998. The watershed science approach (WSA). A methodology developed for the evaluation of geomorphic processes in Bay Area watersheds. San Francisco Estuary Institute, Richmond, California. 31 pp.
- Davis, W. S., and Simons, T.P. (eds), 1996. *Biological assessment and criteria: tools for resource planning and decision making*. Lewis Publishers, Boca Raton, Florida. 432 pp.
- DeVries, P., Burges, S.J., and Daigneau, J., 2001. Measurement of the temporal progression of scour in a pool-riffle sequence in a gravel bed stream using an electronic scour monitor. *Water Resources Research*, v. 37, p. 2805-2816.
- Doberstein, C.P., Karr, J.R., and Conquest, L.L., 2000. The effect of fixed-count sub-sampling on macroinvertebrate biomonitoring in small streams. *Freshwater Biology*, v. 44, p. 355-371.
- Dunne, T. and Leopold, L.B., 1978. *Water in environmental planning*. W.H. Freeman, San Francisco, California. 818 pp.
- Ellen, S.D., 1989. How slides transform into flows: what happens to hillsides in the San Francisco Bay region during and after heavy rains. In: Hodges, C.A., (ed). *Geohazards '88; a symposium highlighting research on the causes, effects, and prediction of geologic and hydrologic hazards*. U.S. Geological Survey, Menlo Park, California. Abstract C 1038, p.12.
- Flosi, G., Downie, S., Hopelain, J., Bird, M., Coey, R., and Collins, B., 1998. *California salmonid stream habitat restoration manual*, 3<sup>rd</sup> edition. State of California, Department of Fish and Game, Inland Fisheries Division.

- Forest, Soil and Water, Inc., O'Connor Environmental, Inc., and East-West Forestry, 1998. Watershed assessment and cooperative in-stream monitoring plan for the Garcia River, Mendocino County, California. Compiled for Mendocino County Resources Conservation District and the California Department of Forestry and Fire Protection, January, 1998.
- Fox, K.F., Jr., Fleck, R.J., Curtis, G.H., and Meyer, C.E., 1985. Implications of the northwestwardly younger age of the volcanic rocks of west-central California. *Geological Society of America Bulletin*, v. 96, p. 647-654.
- Graham, S.A. and Dickinson, W.R., 1978. Apparent offsets of on-land geologic features across the San Gregorio-Hosgri fault trend. In: Silver, E.A., and Normark, W.R., (eds). *San Gregorio-Hosgri fault zone, California*. California Division of Mines and Geology Special Report no. 137, p. 13-22.
- Harrelson, C., Rawlins, C., and Potyondy, J., 1994. Stream channel reference sites: an illustrated guide to field technique. RM-GTR-245, U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.
- Harrington, J.M., 1996. California stream bioassessment procedures. California Department of Fish and Game, Water Pollution Control Laboratory. Rancho Cordova, California. 192 pp.
- Hession, W.C., Pizzuto, J.E., Johnson, T.E., and Horwitz, R.J., 2003. Influence of bank vegetation on channel morphology in rural and urban watersheds. *Geology*, v. 31, p. 147-150.
- Hilton, S. and Lisle, T.E., 1993. Measuring the fraction of pool volume filled with fine sediment. Res. Note PSE-RN-414. Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture, Albany, California. 11 p.
- Huang, H.Q. and Nanson, G.C., 1997. Vegetation and channel variation; a case study of four small streams in southeastern Australia. *Geomorphology*, v. 18, p. 237-249.
- Jayko, A.S., DeMouthe, J., Lajoie, K.R., Ramsey, D.W., and Godt, J.W., 1999. Map showing locations of damaging landslides in San Mateo County, California, resulting from 1997-98 El Niño rainstorms. U.S. Geological Survey Miscellaneous Field Studies map MF-2325H, scale 1:125,000.
- Jones, R.C. and Clark, C.C., 1987. Impact of watershed urbanization on stream insect communities. *Water Resources Bulletin*, v. 23, p. 1047-1055.
- Karr, J.R., 1998. Rivers as sentinels: using the biology of rivers to guide landscape management. In: Naiman, R.J., and Bilby, R.E. (eds). *River ecology and management: lessons from the Pacific coastal ecoregion*. Springer Publishers, New York City, New York. p. 502-528.
- Kobayashi, S. and Kagaya, T., 2002. Differences in litter characteristics and macroinvertebrate assemblages between litter patches in pools and riffles in a headwater stream. *Limnology*, v. 3, p. 37-42.
- Koehler, J.T., 2003. Fish habitat assessment: a component of the watershed management plan for the Carneros Creek watershed, Napa County, California. A technical report of the Napa County Resource Conservation District, Napa, California. 36 pp.
- Kondolf, G.M., 1988. Salmonid spawning gravels: a geomorphic perspective on their distribution, size modification by spawning fish, and application of criteria for gravel quality. PhD thesis, The Johns Hopkins University.

- Kondolf, G.M., 2000. Assessing salmonid spawning gravel quality. *Transactions of the American Fisheries Society*, v. 129, p. 262-281.
- Lammert, M. and Allan, J.D., 1999. Assessing biotic integrity of streams: effects of scale. In: *Measuring the influence of land use/cover and habitat structure on fish and macroinvertebrates*. *Environmental Management*, v. 23, p. 257-270.
- Lenat, D.R. and Crawford, J.K., 1994. Effects of land use on water quality and aquatic biota of three North Carolina Piedmont streams. *Hydrobiologia*, v. 294, p. 185-199.
- Leopold, L.B., 1994. *A View of the River*. Harvard University Press, Cambridge, Massachusetts. 298 pp.
- Leopold, L.B., Wolman, M.G., and Miller, J.P., 1964. *Fluvial Processes in Geomorphology*. W.H. Freeman, New York City, New York. 522 pp.
- Leopold, L.B. and Maddock, T., 1953. The hydraulic geometry of stream channels and some physiographic implications. *U.S. Geological Survey Professional Paper 252*.
- Lewis, P.A., Klemm, D.J., and Thoeny, W.T., 2001. Perspectives on use of a multimetric lake bioassessment integrity index using benthic macroinvertebrates. *Northeastern Naturalist*, v. 8, p. 233-246.
- Li, J., Herlihy, A., Gerth, W., Kaufmann, P., Gregory, S., Urquhart, S., and Larson, D.P., 2001. Variability in stream macroinvertebrates at multiple spatial scales. *Freshwater Biology*, v. 46, p. 87-97.
- Lisle, T.E., 1999. Channel processes and watershed function. In: Taylor, R. N. (ed). *Proceedings of a workshop: using stream geomorphic characteristics as a long-term monitoring tool to assess watershed function*. Humboldt State University, Arcata, California. p. 4-14.
- Lisle, T.E., and Hilton, S., 1999. Fine bed material in pools of natural gravel bed channels. *Water Resources Research*, v. 35, p. 1291-1304.
- Lisle, T.E., Ikeda, H., and Iseya, F., 1991. Formation of stationary alternate bars in a steep channel with mixed-size sediment: a flume experiment. *Earth Surface Processes and Landforms*, v. 16, p. 463-469.
- Lloyd, D.S., 1987. Turbidity as a water quality standard for salmonid habitats in Alaska. *North American Journal of Fisheries Management*, v. 7, p. 33-45.
- McCabe, G.T., Coley, T.C., Emmett, R.L., Muir, W.D., and Durkin, J.T., 1981. The effects of the eruption of Mt. St. Helens on fishes in the Columbia River Estuary. *Estuaries*, v. 4, p. 247.
- McKee, L., Leatherbarrow, J., Newland, S., and Davis, J., 2003. A review of urban runoff processes in the Bay Area: Existing knowledge, conceptual models, and monitoring recommendations. A report prepared for the Sources, Pathways and Loading Workgroup of the Regional Monitoring Program for Trace Substances. San Francisco Estuary Institute, Oakland, California. Contribution no. 66.
- Millar, R.G., 2000. Influence of bank vegetation on alluvial channel patterns. *Water Resources Research*, v. 36, p. 1109-1118.
- Milliken, R., 1995. *A time of little choice: the disintegration of tribal culture in the San Francisco Bay Area 1769-1810*. Ballena Press, Menlo Park, California. 364 p.

- Montgomery, D.R. and Buffington, J.M., 1997. Channel-reach morphology in mountain drainage basins. *Geologic Society of America Bulletin*, v. 109, p. 596-611.
- Newcombe, C.P. and MacDonald, D.D., 1991. Effects of suspended sediments on aquatic ecosystems. *North American Journal of Fisheries Management*, v. 11, p. 72-82.
- Pearce, S., O'Connor, M., Grossinger, R., and McKee, L., 2002. Napa River sediment TMDL baseline study: geomorphic processes and habitat form and function in Soda Creek. San Francisco Estuary Institute, Oakland, California. Contribution no. 63.
- Pearce, S., O'Connor, M., McKee, L., and Jones, B., 2003. Channel geomorphology assessment: A component of the watershed management plan for the Carneros Creek watershed, Napa County, California. A Technical Report of the Regional Watershed Program. San Francisco Estuary Institute, Oakland, California. Contribution no. 67. 58 pp.
- Pearce, S., O'Connor, M., McKee, L., and Jones, B., 2003. Channel geomorphology assessment: A component of the watershed management plan for the Sulphur Creek watershed, Napa County, California. A Technical Report of the Regional Watershed Program. San Francisco Estuary Institute, Oakland, California. Contribution no. 68. 60 pp.
- Plafker, G., and Galloway, J.P. (eds), 1989. Lessons learned from the Loma Prieta, California earthquake of October 17, 1989. U.S. Geological Survey Professional Paper 1045. 48 pp.
- Rantz, S.E., 1971a. Mean annual precipitation depth-duration-frequency data for the San Francisco Bay Region. U.S. Geological Survey, Basic data contribution 32. 24 pp.
- Rantz, S.E., 1971b. Suggested criteria for hydrologic design of storm-drainage facilities in the San Francisco Bay Region California. U.S. Geological Survey Open File Report 3019-11. 72 pp.
- Regional Water Quality Control Board, 1997. San Francisco Bay Basin (Region 2) Water Quality Control Plan. Regional Water Quality Control Board, Oakland California.
- Reisner, D.W. and Bjornn, T.C., 1979. Habitat requirements of anadromous salmonids. In: Meehan, W.R. (ed). Influence of forest and range management on anadromous fish habitat in western America. Pacific Northwest Forest and Range Experiment Station. U.S. Department of Agriculture, Forest Service, Portland, Oregon. General Technical Report, PNW-96.
- Resh, V.H. and Jackson, J.K., 1993. Rapid assessment approaches to biomonitoring using benthic macroinvertebrates. In: Rosenberg, D.M. and Resh, V.H. (eds). Freshwater biomonitoring and benthic macroinvertebrates. Chapman and Hall, New York City, New York. p. 195-233
- Richards, G., 1973. Crossroads: people and events of the redwoods of San Mateo County. Gilbert Richards Publications, Woodside, California. 128 pp.
- Rosgen, D., 1996. Applied river morphology. Wildland Hydrology, Pasoga Springs, Colorado.
- Rosi-Marshall, E. and Wallace, J.B., 2002. Invertebrate food webs along a stream resource gradient. *Freshwater Biology*, v. 47, p.129-141.
- Sarna-Wojcicki, A.M., Lajoie, K.R., Meyer, C.E., Adam, D.P., and Rieck, H.J., 1991. Tephrochronologic correlation of upper Neogene sediments along the Pacific margin, conterminous United States. In: Morrison, R.B. (ed). Quaternary nonglacial geology; conterminous United States. *The Geology of North America*, v. K-2. Geological Society of America, Boulder, Colorado.

- Shirazi, M.A. and Seim, W.K., 1979. A stream system evaluation – an emphasis on spawning habitat for salmonids. EPA-600/3-79-109. U.S. Environmental Protection Agency, Corvallis Environmental Research Laboratory. Corvallis, Or. 44 pp.
- Sigler, J.W., Bjornn, T.C., and Everest, F.H., 1984. Effects of chronic turbidity on density of steelheads and coho salmon. *Transactions of the American Fisheries Society*, v. 113, p. 142-150.
- Simon, A. and Collison, A.J., 2002. Quantifying the mechanical and hydrologic effects of riparian vegetation on streambank stability. *Earth Surface Processes and Landforms*, v. 27, p. 527-546.
- Simpson, G. D., Thompson, C., Noller, J. S., and Lettis, W. R., 1997. The northern San Gregorio fault zone: Evidence for the timing of Late Holocene earthquakes near Seal Cove. *Bulletin of the Seismological Society of America*, v. 87, p. 1158-1170.
- Spence, B.C., Lomnický, G.A., Hughes, R.M., and Novitzki, R.P., 1996. An ecosystem approach to salmonid conservation. *Management Technology*, Corvallis, Oregon. TR-4501-96-6057.
- Stanger, F.M., 1967. Sawmills in the redwoods: Logging on the San Francisco Peninsula 1849-1967. San Mateo County Historical Association, San Mateo, California. 160 pp.
- Strahler, A.N., 1957. Quantitative analysis of watershed geomorphology. *American Geophysical Union Transcripts*, v. 38, p. 913-920.
- Thompson, J.M., and Evernden, J. F., 1986. Map showing predicted seismic-shaking intensities of and earthquake in San Mateo County, California comparable in magnitude to the 1906 San Francisco Earthquake. U.S. Geological Survey Miscellaneous Field Investigations Map I-1257-H, scale 1:62,500.
- Turner, D.L., 1970. Potassium-Argon dating of Pacific Coast Miocene foraminiferal stages. In: Bandy, O.L. (ed). *Radiometric dating and Paleontologic zonation: Geological Society of America Special Paper 124*, p. 91-129.
- United States Geological Survey, 2003. Topographic information: 10 m digital elevation model (DEM): <http://edc.usgs.gov/products/elevation/dem.html>. (June, 2003).
- Usseglio-Polatera, P., Bournaud, M., Richoux, P., and Tachet, H., 2000. Biological and ecological traits of benthic freshwater macroinvertebrates: relationships and definition of groups with similar traits. *Freshwater Biology*, v. 43, p. 175-205.
- Vannote, R.L., Minshall, G.W., Cummins, K.W., Sedell, J.R., and Cushing, C.E., 1980. The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, v. 37, p. 130-137.
- Ventura County Watershed Protection District, 2001. Ventura River watershed biological and physical/habitat assessment, Ventura County, California. Ventura Countywide Stormwater Quality Management Program, Ventura, California. 22 pp.
- Vyverberg, K.A., Snider, B., and Titus, R.G., 1997. Lower American River chinook salmon spawning habitat evaluation: An evaluation of attributes used to define the quality of spawning habitat. California Department of Fish and Game, Sacramento, California.
- Wagner, R.J. and Nelson, R.E., 1954. Soil Survey of the San Mateo Area, California. U.S. Department of Agriculture, Soil Conservation Service, no. 13, issued 1961.

- Washington Forest Practices Board, 1997. Standard methodology for conducting watershed analysis.
- Weber, G.E, 1994. Late Pleistocene slip rates on the San Gregorio fault zone at Point Año Nuevo, San Mateo County, California. In: Lettis, W.R. (ed). Field trip guidebook to transpressional deformation in the San Francisco Bay Area, Friends of the Pleistocene Pacific Southwest Cell. 28 pp.
- Weber, G.E. and Cotton, W.R., 1981. Geologic investigation of recurrence intervals and recency of faulting along the San Gregorio fault zone, San Mateo County, California. U.S. Geological Survey Open File Report OF-81-0257. 131 pp.
- Wieczorek, G.F., 1981. Ground water level and precipitation data for slopes near La Honda, California. U.S. Geological Survey Open File Report OF-81-367. 25 pp.
- Wieczorek, G.F., 1982. Map showing recently-active and dormant landslides near La Honda, central Santa Cruz Mountains, California. U.S. Geological Survey Miscellaneous Field Studies Map MF-1422, scale 1:4,800.
- Wieczorek, G.F., 1984. Preparing a detailed landslide-inventory map for hazard evaluation and reduction. Bulletin of the Association of Engineering Geologists, v. 21, p. 337-342.
- Wieczorek, G.F. and Keefer, D.K., 1984. Landslide at La Honda, California probably triggered by the April 24, 1984 Morgan Hill Earthquake. In: Hoose, S.N., 1984. The Morgan Hill, California earthquake of April 24, 1984: a preliminary report. U.S. Geological Survey Open File Report OF-84-498, p. 109-115.
- Wieczorek, G.F., 1987. Effect of rainfall intensity and duration on debris flows in central Santa Cruz Mountains, California. Geological Society of America Reviews in Engineering Geology, v. 7, p. 93-104.
- Wieczorek, G.F., Brown, W.M., Mark, R.K., Rice, P., and Alger, C.S., 1989. La Honda landslide test area, San Mateo County, California. In: Brown, W.M. (ed). Landslides in Central California. Fieldtrips for the 28<sup>th</sup> International Geologic Congress, guidebook T381. American Geophysical Union, Washington D.C. p. 39-44.
- Williamson, D.A., 1984. Unified Rock Classification System. Bulletin of the Association of Engineering Geologists, v. 21, p. 345-354.
- Wilson, R.C. and Wieczorek, G.F., 1995. Rainfall thresholds for the initiation of debris flows at La Honda, California. Environmental and Engineering Geoscience, v. 1, p. 11-27.
- Wolman, M.G. and Miller, J.P., 1960. Magnitude and frequency of forces in geomorphic processes. Journal of Geology, v. 68, p. 54-74.
- Wooster, J., 2000. Large woody debris volumes and accumulation rates in cleaned streams in redwood forests near the confluence of South and Mainstem Eel River, California. Draft report presented at Large Woody Debris in North Coast Streams Conference, Ukiah, California, May 2003.
- Western Regional Climate Center, 2003. [www.wrcc.dri.edu](http://www.wrcc.dri.edu) (<http://www.wrcc.dri.edu/summary/climsmnca.html>) (April, 2003).
- Yoder, C.O. and Rankin, E.T., 1998. The Role of Biological Indicators in a State Water Quality Management Process. Environmental Monitoring and Assessment, v. 51, p. 61-88.