Rebecca Sutton,¹ Shavonne K. Stanek,¹ Sherri A. Mason,² Ellen Willis-Norton,¹ Ian F. Wren,³ Carolynn Box⁴ San Francisco Estuary Institute, 4911 Central Avenue, Richmond, CA 94804 2 State University of New York at Fredonia, 280 Central Avenue, Science Complex 340, Fredonia, NY 14063 San Francisco Baykeeper, 1736 Franklin Street, Suite 800, Oakland, CA 94612 4 5 Gyres, 3131 Olympic Blvd, Suite 302, Santa Monica, CA 90404

SAN FRANCISCO BAY

FRAGMENTS OF PLASTIC 5 MM OR SMALLER

Sources of aquatic microplastic pollution include microbeads used in personal care products such as facial scrubs and toothpastes, pellets (called nurdles) used as precursors for industrial products, plastic fibers derived from washing clothes made with synthetic materials, and fragments of larger plastic items. Motivated by recent state and federal efforts to ban microbeads in personal care products, the Regional Monitoring Program for Water Quality in San Francisco Bay (RMP) characterized Bay surface waters and wastewater treatment plant (WWTP) effluents for microplastic contaminants. Nine Central and South Bay surface water samples were collected using a Manta Trawl. Two-hour sieved samples of effluent were collected from eight WWTPs discharging to the Bay. Microplastics in samples were characterized by size, type, and abundance. Preliminary results from this survey for plastic pollution in the San Francisco Bay are presented.

METHODS

SAMPLE COLLECTION

EIGHT BAY AREA WWTPS: Treated effluent was sieved through 0.355 mm and 0.125 mm mesh, typically for two hours during peak flow.



30-minute trawl. In some areas, trawl contents included considerable vegetation; nine small fish were collected as accidental by-catch at one site.

Although fish collection was not planned, the abundance of microplastics in these by-catch fish was determined after thorough rinsing to remove external contaminants.



hoto of effluent sample collection

courtesy Eric Dunlavey

Ian Wren of San Francisco Baykeeper deploying the Manta Trawl. hoto courtesy Meg Sedlak

SAMPLE PREPARATION

Surface water samples required DI water to recover associated plastic different sized nets (0.355 mm, 1 mm a wet peroxide oxidation (WPO) process was used to remove natural organic material, leaving behind

The WWTP effluent samples also went



Wet peroxide oxidation (WPO) process: Before (left) and after (right) Photo courtesy Sherri A. Mason

considerable preprocessing. Vegetation was rinsed in triplicate, then soaked in debris. Samples were sieved into three and 4.75 mm), rinsed and categorized. After larger plastic debris was removed,

through the WPO process.



MICROBEADS

made up of small pellets and

tragments used in personal

care products such as tacial

scrubs and toothpastes – see

Focus on Microbeads

NURDLES

precursors to manutacture plastic products

FRAGMENTS

PATHWAYS FOR PLASTICS TO ENTER THE BAY

- Wind- and stormwater-carried trash and plastic debris from land
- Illegal dumping [3]
- Microbeads from personal care products and plastic fibers from clothing treatment is not specifically designed to remove microplastics, so these particles can be released in treated effluent [5]

POTENTIAL CONCERNS OF MICROPLASTICS

- entially sorb to microplastics [6]
- Lower trophic organisms can mistake microplastics for food; ingestion can lead to physical harm, exposure to sorbed contaminants, and bioaccumulation of microplastics in higher trophic organisms [1,6,7]



Microplastics

0 2.5 5 10 Miles

Abundance (Count/Km2

1000001 - 1500000

1500001 - 2000000

14000 - 500000

500001 - 1000000

FOCUS ON MICROBEADS

sign of microbead contamination, these particles are only a small

A number of recent state, federal, and international actions aim to ban or limit the use of plastic microbeads in personal care products. Many companies have already committed to switching to alternate ingredients.

RESULTS

WWTP EFFLUENT

- The eight WWTPs discharged an average of 6,900,000 particles of microplastic per
- The average count of plastic particles per gallon was higher than observed in a similar study of New York state WWTPs (0.33 vs. 0.08 particles per gallon), as was the average discharge per day (6,900,000 vs. 420,000 particles per
- WWTPs employing more advanced wastewater treatment technologies did not have lower concentrations of microplastics than more traditional secondary technology
- Fibers were the dominant form of microplastic pollution in effluent, and are likely derived from washing synthetic clothing and fabrics (FIGURE 1)
- Fragments, the second most abundant form of microplastic pollution, may be derived from microbeads in personal care products as well as other sources
- The absence of small pellets does not indicate a lack of microbead-based contamination, as most microbeads are classified as fragments [2]
- A breakdown of WWTP effluent microplastic abundance by size shows they were fairly evenly split between smaller and larger particles (TABLE 2)
- A 24-hour sample could provide a more comprehensive picture of microplastic pollution in treated wastewater, as peak personal care product use follows distinct diurnal patterns

FILM 2% PELLET 0% Microplastic Microplastic relative particles by type abundance by type FIBER 27% detected i detected in Bay Bay WWTP surface water effluent samples

FRAGMENT (INCLUDES MICROBEADS, OTHER SOURCES) PELLET (MICROBEADS, NURDLES) FIBER (SYNTHETIC FABRIC, CLOTHING, FISHING LINES) FILM (PLASTIC BAGS, PACKAGING)

FOAM (STYROFOAM, CIGARETTE BUTTS)

WWTP	PARTICLES/ GALLON	PARTICLES/ DAY
San Jose-Santa Clara	0.18	15,000,000
East Bay MUD	0.27	12,000,000
Palo Alto	0.48	9,600,000
Central Contra Costa	0.27	8,100,000
Fairfield-Suisun	0.35	4,100,000
EBDA/San Leandro	0.082	4,100,000
San Mateo	0.24	2,000,000
SFO (sanitary plant)	0.74	460,000
Average	0.33 ± 0.19	$6,990,000 \pm 4,700,000$

 TABLE 1. Microplastic levels in Bay

PLASTIC TYPE	0.125 – 0.354 mm	>0.355 mm
Fragment	53	34
Pellet	0	0
Fiber	216	193
Film	1	10
Foam	3	1
Total Count	273	238
% of Total	53%	47%

TABLE 2. WWTP effluent microplastic abundance by type and size (total count)

ABUNDANCE/k (AVERAGE) 1,000,000 Patapsco River Chesapeake Bay[9] Magothy River Chesapeake Bay[9] Rhode River Chesapeake Bay[9] Corsica River Chesapeake Bay[9] Lake Superior[3] Lake Huron[3] 3,000

 TABLE 3. San Francisco Bay has higher aver age levels of microplastic contamination than the Great Lakes or Chesapeake Bay

PLASTIC TYPE	0.355 – 0.999 mm	1.000 – 4.749 mm	>4.75 mm
Fragment	280,000	99,000	3,900
Pellet	11,000	1,800	0
Fiber	110,000	77,000	2,100
Film	18,000	30,000	5,300
Foam	21,000	34,000	1,500
Count/km ²	440,000	240,000	13,000
% of Total	63%	35%	2%

TABLE 4. Average Bay microplastic abundance by type and size (abundance/km²)

BAY SURFACE WATER

- South Bay levels of microplastic contamination are typically higher than Central Bay
- Average South and Central Bay microplastic levels are higher than average measurements from the Great Lakes and Chesapeake Bay
- Fragments were the dominant form of microplastic pollution in surface water, and may be derived from microbeads in personal care products as well as many other sources
- Fibers, the second most abundant form of microplastic pollution, may be derived from fishing line as well as washing synthetic clothing
- Pellets, from microbead products and pre-production plastic nurdles, are a smaller portion of overall Bay surface water microplastic pol-
- Differences in the relative proportion of plastic types in effluent (FIGURE 1) and Bay surface water (FIGURE 3) may be due to sources of plastic from other pollution pathways (e.g., stormwater) or in situ processes such as binding with natural particles, settling to the Bay floor, or ingestion by biota
- A breakdown of Bay microplastic abundance by size shows a greater proportion of smaller (0.355 – 0.999 mm) microplastic particles TABLE 4); smaller particles are more easily ingested by aquatic organisms [1]

- 52 pieces of microplastic were recovered from nine small prey fish collected as unintentional by-catch at one surface water site; two small pellets clearly derived from microbead products
- These Bay fish averaged nearly 6 pieces of microplastic per fish; in contrast, 1-3 pieces of microplastic are typically recovered from fish in the Great Lakes [10]
- These preliminary findings suggest further study of microplastic contamination of Bay fish is needed to determine whether Bay fish contain more microplastics than those in other major water bodies, and to investigate the potential for bioaccumulation of microplastics in sport fish consumed by people

CONCLUSIONS

MICROPLASTICS ARE WIDESPREAD in the Central and South Bays, and found at levels higher than other water bodies near highly urbanized regions of the U.S.

BAY WWTPs discharge levels than WWTPs in

THE DATA DO NOT SUGGEST a difference in the concentration of microplastics in effluent for WWTPs employing secondary vs. advanced secondary treatment

FRAGMENTS AND

greatest abundance in both Bay surface water and effluent

Microplastic particles recovered from the first rinse of

vegetation from a surface

water trawl sample. Photo

courtesy Sherri A. Mason

ANALYSIS

Plastic particles were

typed and counted.

flow rates (WWTP

samples) or tow

abundances.

Éxtrapolations using

lengths (surface water

samples) were used to

calculate microplastics

MICROBEADS IN PERSONAL CARE

PRODUCTS, a recent policy focus, consist primarily of small fragments, and to a lesser extent small pellets; our findings indicate microbeads can be found in the Bay, and are other regions likely discharged via treated wastewater

MONITORING MICROPLASTICS

in Bay sediment and biota will provide a more complete picture of Bay contamination relative to

ACKNOWLEDGMENTS

Funding for this study was provided by the Regional Monitoring Program for Water Quality in San Francisco Bay, with in-kind contributions from San Francisco Baykeeper and 5 Gyres. Special thanks to the WWTPs that voluntarily participated in this study, and to the staff that made sample collection possible: T. Nguyen and M. LaBella (Central Contra Costa); M. Connor, S. Abbasi and A. Berumen (EBDA/San Leandro); N. Arsem and J. Allard (EBMUD); G. Heravian, M. Herston, B. Carver and T. Harshell (Fairfield-Suisun); K. North (Palo Alto); B. Zolan (SFO); E. Dunlavey and J. Ervin (San Jose-Santa Clara); P. Dalla-Betta and S. Scheidt (San Mateo). Additional thanks to Oberlii College, Career Center for providing intern support. Poster design by Linda Wanczyk.

REFERENCES

in Bay surface water

Farley, H., Amato, S. 2013. Microplastic pollution in the surface water of the Laurentian Great Lakes. Marine Pollution Bulletin 77, 177-182

FIGURE 2. Total microplastic abundances

- [4] Browne, M.A., Crump, P., Niven, S.J. et al. 2011. Accumulation microplastic on shorelines worldwide: Sources and sinks. Environmental Science & Technology. 45, 9175-9179.
- 5] Hogue, C. 2013. "Microplastic Beads Pollute Great Lakes." Chemical and Engineering News 16 Sept. 2013, 91st ed.: 23-25. 6] Seltenrich, N. 2015. New Link in the food chain? Marine plastic pollution and seafood safety. Environmental Health Perspectives

microplastics by zooplankton in the North Pacific Ocean. Archives of

Mason, S.A., Ricotta, R., Smith, M.L., Vinson, M. Microplastic

contaminates within the Laurentian Great Lakes food web' intended

licro-plastic Pollution: A Comparative Survey of Wastewater Effluen New York. Center for Earth and Environmental Science Student ttp://digitalcommons.plattsburgh.edu/cees_student_posters/8 9] Yonkos L. T., Friedel, E. A., Perez-Reyes, A. C., Ghosal, S., Arthur, C. 2014. Microplastics in four estuarine rivers in the Chesapeake Bay, U.S.A. Environmental Science & Technology 48, 14195-14202

AQUATIC SCIENCE CENTER 4911 Central Ave, Richmond, CA 94804, 510-746-7334, www.sfei.org

CONTACT INFORMATION

Rebecca Sutton, Ph.D. – RebeccaS@sfei.org



