Introduction

Historically, the San Francisco Estuary has been considered a low primary productivity estuary, despite having relatively high concentrations of nutrients. The low primary production has been attributed to the Estuary being light-limited due to high suspended sediment concentrations and the high energy of wind-wave action. However, recent changes in phytoplankton community structure and Bay ecology, decreases in dissolved oxygen, and a decrease in suspended sediment concentrations have raised concerns about the Estuary's ability to continue to be resilient to the impacts of continuing nutrient loads.



The Regional Monitoring Program for Water Quality in the San Francisco Estuary (RMP) has monitored nutrients using a targeted sampling design since 1993, and beginning in 2002, a probabilistic sampling design employing the US EPA's Generalized Random Tessellation Stratified (GRTS) sample design. Dissolved nitrate and phosphate concentrations from samples collected at the probabilistic stations (2002-2009) were investigated for spatial distributions, while concentrations from samples collected at five fixed historic stations (1993-2009) were examined for temporal trends (Figure 1).

RMP water sampling map showing fixed

historical stations (1993-2009), and Gen-

SanPablo

Strait

RMP Water Stations

Bay Bathymetry

0 to 6

Depth Below Sea Level (ft)

GRTS Random Stations

Historic Fixed Stations

eralized Random Tessellation Stratified

(GRTS) random stations (2002-2009).

hymetry data from the CaDFG, 2002

lap Created by Marcus Klatt, SFEI in

AD 1983 California Teale Albers.



DISSOLVED NITRATE AND PHOSPHATE CONCENTRATIONS IN SAN FRANCISCO ESTUARY SURFACE WATERS: SPATIAL DISTRIBUTION (2002-2009) AND TEMPORAL TRENDS (1993-2009)

John R.M. Ross, San Francisco Estuary Institute, 7770 Pardee Lane, 2nd Floor, Oakland, CA 94621-1424 (john@sfei.org)

Methods

SPATIAL COMPARISONS

GRTS sites are allocated into five hydrographic regions of the Estuary: Suisun Bay, San Pablo Bay, Central Bay, South Bay, and Lower South Bay. Regional means, standard deviations, and cumulative distribution functions (CDFs) were estimated using the R statistical software environment rg/), and version 2.2 of the spsurvey statistical liwas designed by the US EPA specifically for the statistical analysis of GRTS spatial survey designs.

Differences between regional CDFs were evaluated using the F distribution version of the Wald statistic, with nondetects replaced with one-half the method detection limit. Substitution, however, can produce an invasive pattern alien to the concentrations actually in the samples, resulting in generally poor estimates and incorrect statistical tests (Helsel, 2005; Helsel 2009). Therefore, differences between the dissolved nitrogen cumulative distribution functions were also investigated using a Generalized Wilcoxon (Score) Test for censored data followed by multiple pairwise comparisons (Helsel, 2005). Comparisons were considered significant at a Bonferroni corrected alpha of 0.005.

Pairwise Comparison

TEMPORAL TRENDS

At each station without nondetects trends were examined by ordinary least squares (OLS) linear regression analysis using untransformed concentrations as the dependent variable, and sampling date as the independent variable. Normality of the residuals was evaluated using the Anderson-Darling test, and when sufficient evidence was observed to reject the null hypothesis of normality (p<0.10), attempts were made to normalize by transforming the data. If data transformation was unsuccessful, then a robust regression analysis was conducted using an M-estimation robust regression technique called iteratively reweighted least squares (Chatterjee and Machler, 1997). The presence of first-order autocorrelation was examined using the Durbin-Watson test, but no conclusive evidence of firstorder autocorrelation was found in the data.

Trends at each station with nondetects were evaluated by the Akritas-Thiel-Sen method for censored data (Helsel, 2005) using untransformed concentrations as the dependent variable, and sampling date as the independent variable. Unusual observation, observations with standardized residuals >2 or <-2, were identified in the majority of the regressions, but lacking a reason for exclusion were considered extreme, but valid measurements (US EPA, 1992). A significant positive slope (p<0.05) was assumed to indicate an increase at the station over time. Similarly, a significant negative slope assumes a decrease over time, while a lack of significance indicates no change in water concentration over time.

Table 1 Results of multiple pairwise comparisons between regional cumulative distribution functions (CDFs).

	Central Bay vs. Lower South Bay	6.84	2	66	0.002	0.005	Yes
	Central Bay vs. South Bay	4.71	2	80	0.012	0.005	No
	Central Bay vs. San Pablo Bay	3.76	2	54	0.030	0.005	No
	Central Bay vs. Suisun Bay	4.15	2 2 2	53 93 67 66 81	0.021	0.005 0.005 0.005 0.005 0.005	No Yes No
	Lower South Bay vs. South Bay	9.53 0.70 2.86 9.02			0.000		
	Lower South Bay vs. San Pablo Bay				0.498		
	Lower South Bay vs. Suisun Bay		2		0.064		No
	South Bay vs. San Pablo Bay		2		0.000		Yes
	South Bay vs. Suisun Bay	14.39	2	80	0.000	0.005	Yes
	San Pablo Bay vs. Suisun Bay	0.74	2	54	0.484	0.005	No
	Generalized Wilcoxon (Score) Test	Chi-Square	df		р		
		30.5923	4		0.000		
	Pairwise Comparison						
	Central Bay vs. Lower South Bay	1.60	1		0.206	0.005	No
	Central Bay vs. South Bay	8.31	1		0.004	0.005	Yes
	Central Bay vs. San Pablo Bay	2.84	1		0.092	0.005	No
	Central Bay vs. Suisun Bay	6.95	1		0.008	0.005	No
	Lower South Bay vs. South Bay	12.81	1		0.000	0.005	Yes
	Lower South Bay vs. San Pablo Bay	0.06	1		0.812	0.005	No
	Lower South Bay vs. Suisun Bay	0.75	1		0.386	0.005	No
	South Bay vs. San Pablo Bay	19.36	1		0.000	0.005	Yes
	South Bay vs. Suisun Bay	27.70	1		0.000	0.005	Yes
	San Pablo Bay vs. Suisun Bay	0.54	1		0.461	0.005	No
Phosphate							
	Central Bay vs. Lower South Bay	211.02	2	66	0.000	0.005	Yes
	Central Bay vs. South Bay	49.92	2	80	0.000	0.005	Yes
	Central Bay vs. San Pablo Bay	0.32	2	54	0.730	0.005	No
	Central Bay vs. Suisun Bay	4.32	2	53	0.020	0.005	No
	Lower South Bay vs. South Bay	52.81	2	93	0.000	0.005	Yes
	Lower South Bay vs. San Pablo Bay	85.70	2	67	0.000	0.005	Yes
	Lower South Bay vs. Suisun Bay	277.17	2	66		0.005	Yes
	South Bay vs. San Pablo Bay	18.66	2	81	0.000	0.005	Yes
		108.80	2	80	0.000	0.005	Yes
	South Bay vs. Suisun Bay	100.00	_				

Wald_F df1 df2 p Bonferroni p Significant

Results

Considerable variation was observed in the concentrations of dissolved and phosphate concentrations among the five estuary regions (Figure 2 and Figure 3). Mean dissolved nitrate concentrations in water were spatially distributed as Suisun Bay (0.324, sd = 0.131 mg/l), Lower South Bay (0.311, sd = 0.254 mg/l), San Pablo Bay (0.296, sd = 0.147mg/l), Central Bay (0.235, sd = 0.101 mg/l), and South Bay (0.158, sd = 0.101 mg/l) 0.126 mg/l), and dissolved phosphate concentrations as Lower South Bay (0.348, sd = 0.102 mg/l), South Bay (0.185, sd = 0.053 mg/l), San Pablo Bay (0.126, sd = 0.070 mg/l), Central Bay (0.116, sd = 0.061 mg/l), and Suisun Bay (0.097, sd = 0.038 mg/l).

Significant differences were found between estimated regional CDFs for both dissolved nitrate and phosphate (Table 1). Dissolved nitrate concentrations were significantly lower in the South Bay compared to all regions except the Central Bay, while Lower South Bay concentrations were significantly higher than the Central Bay (Table 1, Figure 4). Pairwise comparisons after a significant Generalized Wilcoxon (Score) Test for censored data found significantly lower concentrations in the South Bay compared to all the other regions (Table 1). The difference in results due to the treatment of nondetects measured in several Central Bay, South Bay, and Lower South Bay water samples. Dissolved phosphate concentrations were significantly greater in the Lower South Bay compared to all the other regions, while concentrations in the South Bay were significantly greater than in San Pablo Bay, Central Bay, and Suisun Bay (Table 1, Figure 5).



Analysis of temporal trends showed a significant decrease in dissolved nitrate concentrations at the Dumbarton Bridge (BA30: South Bay), and Golden Gate (BC20: Reference Site) stations (Table 2). No significant trends were found in dissolved phosphate concentrations.

San Francisco Estuary generated by ordinary measured at Generalized **Random Tessellation** Stratified (GRTS) random stations (2002-2009).

Spatial distribution of

San Francisco Estuary

generated by ordinary

Random Tessellation


stations (2002-2009).

dissolved phosphate in the


kriging of concentrations

measured at Generalized

Stratified (GRTS) random



phosphate in the San Francisco Estuary, CA



— Central Bay – Lower South Bay - South Bay - - San Pablo Bay - Suisun Bay 0.0 0.1 0.2 0.3 0.4 0.5 0.6

Discussion

The bimodal spatial distribution of dissolved nitrate and dissolved phosphate concentrations is consistent with the observations that the northern part of the estuary is strongly influenced by agricultural inputs from the Sacramento and San Joaquin rivers (Cloern et al., 2007), while the southern region, being situated in a densely populated urban watershed, receives 98% of its nitrogen input from municipal wastewater treatment plants (Smith and Hollibaugh, 2006).

The significant decline in dissolved nitrate at the Dumbarton Bridge (BA30: South Bay) station is in agreement with the large declines in dissolved inorganic nitrogen and dissolved reactive phosphorus reported by Cloern et al. (2007) in South San Francisco Bay for the period from January 1990 through December 2005. The nutrient declines are consistent with progressive improvements in municipal wastewater treatment leading to decreasing nitrogen and phosphorous input from the urban watershed (Cloern et al., 2007).



			Nondetects	Regression Type	Transformation	Slope	Intercept	n	р	adj-r²	Anderson-Darling (p value)	Durbin-Watson		Unusual Observations
		StationName					·		•			d	Interpretation	1
ers	BG20	Sacramento River	No	OLS	None	-0.000001	0.300	34	0.876	0.0%	0.514	1.8828	Accept null	Yes
ers	BG30	San Joaquin River	Yes (1)	A-T-S	None	-0.000014	0.779	33	0.394	na	na	na	na	na
ntral Bay	BC10	Yerba Buena Island	No	OLS	None	-0.000004	0.379	32	0.700	0.0%	0.581	2.2421	Accept null	Yes
uth Bay	BA30	Dumbarton Bridge	No	OLS	ln(x)	-0.000059	2.56	33	0.003 ^a	22.7%	0.143	2.4269	Accept null	Yes
ference Site	BC20	Golden Gate	No	OLS	None	-0.000018	0.789	28	0.041 ^a	11.8%	0.882	2.0132	Accept null	Yes
ers	BG20	Sacramento River	No	Robust	None	0.000003	-0.050	34	0.090	5.9%	na	1.6911	Accept null	Yes
ers	BG30	San Joaquin River	No	Robust	None	-0.000002	0.157	33	0.282	0.6%	na	2.2490	Accept null	Yes
ntral Bay	BC10	Yerba Buena Island	No	Robust	None	-0.000002	0.165	33	0.224	1.7%	na	2.2605	Accept null	Yes
uth Bay	BA30	Dumbarton Bridge	No	Robust	None	-0.000011	0.657	33	0.131	4.2%	na	2.3932	Accept null	Yes
ference Site	BC20	Golden Gate	No	Robust	None	-0.000002	0.127	28	0.187	3.0%	na	2.1555	Accept null	Yes
rei ut fe rei nt	ral Bay h Bay rence Site rs rs ral Bay h Bay	rs BG30 ral Bay BC10 h Bay BA30 rence Site BC20 rs BG30 ral Bay BC10 h Bay BA30	rs BG30 San Joaquin River ral Bay BC10 Yerba Buena Island h Bay BA30 Dumbarton Bridge rence Site BC20 Golden Gate rs BG30 Sacramento River rs BG30 San Joaquin River ral Bay BC10 Yerba Buena Island h Bay BA30 Dumbarton Bridge	rs BG30 San Joaquin River Yes (1) ral Bay BC10 Yerba Buena Island No h Bay BA30 Dumbarton Bridge No rence Site BC20 Golden Gate No rs BG30 Sacramento River No rs BG30 San Joaquin River No ral Bay BC10 Yerba Buena Island No h Bay BA30 Dumbarton Bridge No	rs BG30 San Joaquin River Yes (1) A-T-S ral Bay BC10 Yerba Buena Island No OLS h Bay BA30 Dumbarton Bridge No OLS rence Site BC20 Golden Gate No OLS rs BG30 Sacramento River No Robust rs BG30 San Joaquin River No Robust ral Bay BC10 Yerba Buena Island No Robust h Bay BA30 Dumbarton Bridge No Robust	rs BG30 San Joaquin River Yes (1) A-T-S None ral Bay BC10 Yerba Buena Island No OLS None h Bay BA30 Dumbarton Bridge No OLS In(x) rence Site BC20 Golden Gate No OLS None rs BG30 Sacramento River No Robust None rs BG30 San Joaquin River No Robust None ral Bay BC10 Yerba Buena Island No Robust None h Bay BA30 Dumbarton Bridge No Robust None	San Joaquin River Yes (1) A-T-S None -0.000014 ral Bay BC10 Yerba Buena Island No OLS None -0.000004 h Bay BA30 Dumbarton Bridge No OLS In(x) -0.000059 rence Site BC20 Golden Gate No OLS None -0.000018 rs BG20 Sacramento River No Robust None 0.000003 rs BG30 San Joaquin River No Robust None -0.000002 ral Bay BC10 Yerba Buena Island No Robust None -0.000002 h Bay BA30 Dumbarton Bridge No Robust None -0.000011	San Joaquin River Yes (1) A-T-S None -0.000014 0.779 ral Bay BC10 Yerba Buena Island No OLS None -0.000004 0.379 h Bay BA30 Dumbarton Bridge No OLS In(x) -0.000059 2.56 rence Site BC20 Golden Gate No OLS None -0.000018 0.789 res BG20 Sacramento River No Robust None 0.000003 -0.050 res BG30 San Joaquin River No Robust None -0.000002 0.157 ral Bay BC10 Yerba Buena Island No Robust None -0.000002 0.165 h Bay BA30 Dumbarton Bridge No Robust None -0.000011 0.657 Robust Robust None -0.000011 0.657 Robust Robust	San Joaquin River Yes (1) A-T-S None -0.000014 0.779 33 ral Bay BC10 Yerba Buena Island No OLS None -0.000004 0.379 32 h Bay BA30 Dumbarton Bridge No OLS In(x) -0.000059 2.56 33 rence Site BC20 Golden Gate No OLS None -0.000018 0.789 28 res BG20 Sacramento River No Robust None 0.000003 -0.050 34 res BG30 San Joaquin River No Robust None -0.000002 0.157 33 ral Bay BC10 Yerba Buena Island No Robust None -0.000002 0.165 33 h Bay BA30 Dumbarton Bridge No Robust None -0.000011 0.657 33 Robust Robu	San Joaquin River Yes (1) A-T-S None -0.000014 0.779 33 0.394 ral Bay BC10 Yerba Buena Island No OLS None -0.000004 0.379 32 0.700 h Bay BA30 Dumbarton Bridge No OLS In(x) -0.000059 2.56 33 0.003	San Joaquin River Yes (1) A-T-S None -0.000014 0.779 33 0.394 na	San Joaquin River Yes (1) A-T-S None -0.000014 0.779 33 0.394 na na na ral Bay BC10 Yerba Buena Island No OLS None -0.000004 0.379 32 0.700 0.0% 0.581	San Joaquin River Yes (1) A-T-S None -0.000014 0.779 33 0.394 na na na na ral Bay BC10 Yerba Buena Island No OLS None -0.000004 0.379 32 0.700 0.0% 0.581 2.2421	Sam Joaquin River Yes (1) A-T-S None -0.000014 0.779 33 0.394 na na na na na na na n

Literature Cited

Chatterjee, S. and M, Machler, 1997. Robust regression: a weighted least squares approach. Communications in Statistics, Theory and Methods 26, 1381–1394.

Cloern, J.E., Alan D. Jassby, Janet K. Thompson, and Kathryn A. Hieb, 2007. A old phase of the East Pacific triggers new phytoplankton blooms in San Francisco Bay. Proceedings of the National Academy of Sciences of the United States of America 104: 47, 18561-18565

sored environmental data. Hoboken (NJ): John Wiley & Sons,

nondetects in science. The Annals of Occupational Hygiene 54,

exchanges in San Francisco Bay. Limnology and Oceanography 51, 504-517

Helsel, D.R., 2009. Much ado about next to nothing: incorporating United States Environmental Protection Agency (US EPA), 1992 Statistical Training Course for Ground-Water Monitoring Data Analysis, EPA/530-R-93-003, Office of Solid Waste, Washington, for poster design and production and Marcus Klatt for creating the water station sampling map.